

## LIST OF TABLES

Table No.	Title of table	Pg. No
Table 1.1.1	Deaths from Noncommunicable Diseases in India [2016] (Percentage)	34
Table 1.1.2	Targets and Indicators for NCD prevention and control in India under National NCD Monitoring Framework	37
Table 2.4.1.1	List of survey implementing agencies, allotted states and number of PSUs covered for NNMS 2017-18	58
Table 2.7.1	Key quality assurance mechanisms followed	66
Table 4.1.3.1	Occupational status of adults by area of residence and gender (Percentage)	77
Table 4.2.1.1	Current daily tobacco use by area of residence and gender (Percentage)	81
Table 4.2.1.2	Daily tobacco use (any form) by type of product, area of residence and gender (Percentage)	82
Table 4.2.1.3	Number of tobacco products of different types used daily by area of residence and gender (Mean)	82
Table 4.2.1.4	Quit attempts/advices by doctor/health care worker on tobacco use by area of residence and gender (Percentage)	84
Table 4.2.2.1	Patterns of alcohol use in the past 12 months daily or almost daily by area of residence and gender (Percentage)	86
Table 4.2.2.2	Number of standard drinks consumed in one drinking occasion in the past 30 days by area of residence and gender (Mean)	86
Table 4.2.2.3	Maximum number of standard drinks consumed in one drinking occasion by area of residence and gender (Percentage)	87
Table 4.2.2.4	Consumption of alcohol from unauthorized sources in past 7 days by area of residence and gender (Percentage)	88
Table 4.2.3.1	Number of servings of fruits, vegetables and fruit and/or vegetable juices consumed per day by area of residence and gender (Mean)	90
Table 4.2.3.2	Perception related to salt intake by area of residence and gender (Percentage)	91
Table 4.2.3.3	Number of meals consumed outside home in a week by area of residence and gender (Mean)	93
Table 4.2.4.1	Time (minutes) spent in vigorous and moderate level activities per day by area of residence and gender (Mean)	95
Table 4.2.4.2	Time (minutes) spent in physical activity at work, during travel and leisure by area of residence and gender (Mean)	95
Table 4.2.4.3	Voluntary physical activity during recreational time by area of residence and gender (Percentage)	96

Table 4.2.4.4	Time (minutes) spent being sedentary per day by area of residence and gender (Mean)	96
Table 4.2.5.1	Measurements of height, weight, BMI and waist circumference by area of residence and gender (Mean)	97
Table 4.2.5.2	Adults categorized as overweight (including obesity) and obese by area of residence and gender (Percentage)	99
Table 4.2.5.3	Adults with central obesity by area of residence and gender (Percentage)	99
Table 4.2.5.4	Blood pressure measurements by area of residence and gender (Mean)	100
Table 4.2.5.5	Adults with raised blood pressure (known and newly detected) by area of residence and gender (Percentage)	100
Table 4.2.5.6	Blood pressure categories among those measured by area of residence and gender (Percentage)	101
Table 4.2.6.1	Fasting blood glucose levels (mg/dl) by area of residence and gender (Mean)	102
Table 4.2.6.2	Adults with raised fasting blood glucose (known and newly detected) by area of residence and gender (Percentage)	102
Table 4.2.6.3	Fasting blood glucose categories among those measured by area of residence and gender (Percentage)	103
Table 4.2.6.4	Spot urinary Sodium, Potassium and Creatinine excretion levels by area of residence and gender (Mean)	103
Table 4.2.7.1	Adults (40–69 years) with 10-year CVD risk (as per WHO guidelines) by area of residence and gender (Percentage)	105
Table 4.3.1.1	Tobacco use of any form among adolescents by area of residence and gender (Percentage)	108
Table 4.3.1.2	Smoked tobacco use among adolescents by area of residence and gender (Percentage)	108
Table 4.3.1.3	Smokeless tobacco use among adolescents by area of residence and gender (Percentage)	108
Table 4.3.1.4	Current daily tobacco use among adolescents by area of residence and gender (Percentage)	109
Table 4.3.1.5	Adolescents who thought that inhaling smoke from other people's tobacco smoking can cause harm by area of residence and gender (Percentage)	109
Table 4.3.2.1	Age (in years) of initiation of alcohol use among adolescents by area of residence and gender (Mean)	112
Table 4.3.2.2	Source from where alcohol was consumed in past 30 days by area of residence and gender (Percentage)	112

Table 4.3.3.1	Number of days breakfast was skipped in past 30 days by area of residence and gender (Mean)	113
Table 4.3.4.1	Time (minutes) spent in physical activity per day at school by area of residence and gender (Mean)	116
Table 4.3.4.2	Time (minutes) spent in physical activity per day by area of residence and gender (Mean)	116
Table 4.3.4.3	Time (minutes) spent being sedentary in a day by area of residence and gender (Mean)	116
Table 4.3.5.1	Measurements of height, weight and BMI by area of residence and gender (Mean)	117
Table 4.3.5.2	Adolescents categorized as overweight (including obesity) and obese by area of residence and gender (Percentage)	117
Table 4.3.6.1	Reported presence of school/college canteen by area of residence and gender (Percentage)	118
Table 4.3.6.2	Reported availability of food items in school/college canteen by area of residence and gender (Percentage)	118
Table 4.3.6.3	Health promotion and education related information by area of residence and gender (Percentage)	119
Table 4.4.1.1	Source of current consultation and treatment among those with known raised blood glucose by area of residence and gender (Percentage)	123
Table 4.4.2.1	Source of current consultation and treatment among those with known raised blood pressure by area of residence and gender (Percentage)	126
Table 4.4.3.1	Practices reported for blood cholesterol measurement ever in life and history of raised cholesterol by area of residence and gender (Percentage)	127
Table 4.4.3.2	Source of current consultation, treatment and medication for raised blood cholesterol by area of residence and gender (Percentage)	127
Table 4.4.4.1	Adults aged 30–69 years with known cardiovascular conditions and the source of diagnosis by area of residence and gender (Percentage)	129
Table 4.4.4.2	Adults aged 30–69 years who were on medication to prevent or treat known cardiovascular conditions among those with CVDs by area of residence and gender (Percentage)	129
Table 4.4.5.1	Adults aged 30–69 years who had ever undergone oral cancer screening by area of residence, gender and age categories (Percentage)	130
Table 4.4.5.2	Women aged 30–69 years who had ever undergone clinical breast cancer screening by area of residence and age categories (Percentage)	130
Table 4.4.5.3	Women aged 30–69 years who had ever undergone cervical cancer screening by area of residence and age categories (Percentage)	131

Table 4.5.1.1	Overall facilities and infrastructure available in public primary care facilities for all conditions (Percentage)	134
Table 4.5.1.2	Services available for NCDs in public primary care facilities (Percentage)	135
Table 4.5.1.3	Average number of patients with major NCDs attending public primary care facilities in the month prior to the survey	135
Table 4.5.1.4	Services available for management of specific NCDs in public primary care facilities (Percentage)	135
Table 4.5.1.5	Available human resources (medical/paramedical/other staff) in public primary care facilities (Percentage)	136
Table 4.5.1.6	Availability of essential technologies and medicines for NCDs in public primary care facilities (Percentage)	137
Table 4.5.2.1	Overall facilities and infrastructure available in CHCs and district hospitals for all conditions (Percentage)	138
Table 4.5.2.2	Services available for NCDs in CHCs and district hospitals (Percentage)	139
Table 4.5.2.3	Average number of patients attending public secondary health centres in the month prior to the survey	139
Table 4.5.2.4	Service availability for management of NCDs in public secondary health centres (Percentage)	140
Table 4.5.2.5	Available human resources (medical staff) in public secondary health centres (Percentage)	140
Table 4.5.2.6	Available human resources (paramedical/other staff) in public secondary health centres (Percentage)	141
Table 4.5.2.7	Availability of essential technologies and medicines for NCDs as per NPCDCS guidelines in public secondary health centres (Percentage)	142
Table 4.5.3.1	Overall facilities and infrastructure available in private primary health care facilities (Percentage)	144
Table 4.5.3.2	Service availability for management of NCDs in private primary health care facilities (Percentage)	145
Table 4.5.3.3	Availability of essential technologies and medicines for NCDs in private primary health care facilities (Percentage)	145
Table 4.6.1	Time (minutes) spent in practicing asana, pranayam and meditation in a day among those who practiced yoga by area of residence and gender (Mean)	149